

QuralisTM

Driving scientific breakthroughs into
powerful precision medicines for ALS
and other neurodegenerative diseases

May 2025

Driving scientific breakthroughs into powerful precision medicines




Groundbreaking science

Next-gen precision medicines developed by **human disease models** pioneered by QurAlis founders

Targeting **RNA restoration** in validated genetic disease resulting from **mis-splicing targets** in neurodegeneration and beyond

Utilizing **biomarkers** for patient selection, target engagement, and efficacy



First & best-in-class programs

Multiple programs in the clinic with near-term data readouts

Three first-in-class programs for sporadic ALS **on novel genetic targets**

Proprietary FlexASO® platform can potential expand to additional RNA restoration therapies










World-class team to execute

Experienced executive team

UNC13A partnership with Lilly underscores value of FlexASO® platform

\$143.5M equity raised, in addition to Lilly partnership upfront

Advancing precision neuroscience pipeline across multiple clinical and preclinical programs

Program	MOA	Indication	Upcoming Milestone(s)	Discovery	IND-Enabling	Ph1 Safety / PoM ¹	Proof of Concept	Registration Studies	Partner
QRL-201	STMN2	ALS	ANQUR readout H1 2026						
QRL-101	Excitatory (Kv7)	ALS	Patient PoM ¹ study topline Q2/Q3 2025						
		Epilepsy	Proof of Concept initiation YE 2025						
QRL-203	STMN2	FTD (Non-tau)	IND 2026						
QRL-204	UNC13A	ALS / FTD	IND-enabling studies ongoing ²						
QRL-TBA	Undis.	Fragile X	DC ³ 2026						
QRL-TBA	Undis.	PSP	DC 2026						

Lilly

1. PoM = Proof of Mechanism; 2. In partnership with Lilly, clinical plan is to be determined; 3. DC = development candidate nomination

QurAlis' expertise and technologies enable two distinct franchises

Pursuing treatment for CNS disorders with innovative biology and proven modalities

Ion Channel Recovery

(small molecule)

- Neurological disorders often result from ion channel dysfunction
- Kv7.2/7.3 potassium channel is a drug target for >10 high unmet need indications, multiple indications with clinical validation, including:
 - >50% of ALS
 - Epilepsy
 - Pain
 - Mood disorders
- Highly selective Kv7.2/7.3 opener well positioned as potential best-in-class therapeutic:
 - High selectivity, lack of off-target engagement controls AE rates
 - Formulations optimized for different indications

RNA Restoration

(antisense oligonucleotide ,“ASO”)

- Potential to develop first-in-class and best-in-class medicines through FlexASO® platform
 - Active antisense oligonucleotide (ASO) candidates in Phase 1 (1x) and FIH-enabling studies (2x)
- Specifically addresses mis-splicing targets which underly biology of neurodegenerative diseases including:
 - TDP-43-opathies
 - Tau-opathies
 - Fragile X syndrome
- Multiple candidates generated to date with reproducible path to IND and Proof of Concept (PoC)
 - Includes QRL-204 (UNC13A) program licensed to Eli Lilly

Pioneers with unrelenting commitment to patients



Kasper Roet,
PhD
CEO
Co-founder



Emma Bowden,
PhD
Head of
Development



Jason Brown,
MBA
CFO



Hagen Cramer,
PhD
CTO



Dan Elbaum,
PhD
CSO



Vikas Sharma,
PhD
CBO




Robin Wojcieszek,
R.Ph.
Head of
Regulatory Affairs
& Drug Safety






Advised by leading clinicians and scientists in neurodegeneration field

CAB





Angela Genge
MB, FRCP








Ammar Al-Chalabi
MB, ChB, PhD







Merit Cudkowicz
MD, MSc







Dame Pamela J. Shaw
DBE, MBBS, MD,
FRCP, FMedSci,
FAAN, FANA, FAAS







Philip Van Damme
MD, PhD








Leonard van den Berg
MD, PhD








Amy Chappell
MD, FAAN




SAB






Clifford Woolf
MD, PhD








Kevin Eggan
PhD








Matthew Kiernan
MD, PhD








Bernhardt Trout
PhD







Brian Wainger
MD, PhD








Christopher Shaw
MD, PhD






Sudhir Agrawal
PhD





Michelle Hastings
PhD



QA™

© 2025 QURALIS CORP

6

Supported and recognized by investors, pharma, and industry



Investors



EQT
Life Sciences

sanofi ventures



DROIA ventures



mission
BIOCAPITAL



Dementia
Discovery
Fund

inkef capital

ALS
INVESTMENT FUND



MP Healthcare Venture Management, Inc.
Mitsubishi Tanabe Pharma Group



ALEXANDRIA



Awards



FIERCE 15
2020 WINNER



THE
TERMEER
FOUNDATION



Collaborators



QRL-101 in-license
QRL-204 out-license



UMass Chan
MEDICAL SCHOOL

Fragile X

Ion Channel Recovery

Ion channel dysfunction is implicated across wide range of CNS disorders

Kv7.2/3 channel openers have validation across variety of diseases

- GSK's ezogabine was studied in multiple indications including pain, seizure, and mood disorders and marketed for partial-onset seizures before being withdrawn (2017) for undesirable side effect profile, limiting commercial potential
 - Ezogabine also demonstrated signal of disease modification in PoM trial in ALS¹, where hyperexcitability is a key characteristic in up to 40-70% of ALS patients
- Further validation of Kv7.2/3 has been demonstrated by XEN1101 and other clinical programs in epilepsy studies




- QurAlis is developing QRL-101, a highly selective Kv7.2/3 channel opener for ALS and epilepsy with Phase 2/PoC studies initiating by YE'25
 - High affinity to Kv7.2/3
 - Lack of affinity for GABA-A receptors and other Kv7 subtypes
 - Human target engagement observed in Phase 1 studies

Wainger BJ, Macklin EA, Vucic S, et al. *JAMA Neurol.* 2021;78(2):186–196. doi:10.1001/jamaneurol.2020.4300


Kv7 is a clinically validated target in ALS

Ezogabine published trial results¹ (n= 65 patients) validated the importance of reducing hyperexcitability through Kv7

Statistically significant² dose-dependent effects on biomarkers that predict patient survival



Statistically significant² impact on critical disease progression & efficacy biomarker (CMAP)




Nearly all (97%) participants in the trial reported at least one adverse event
Most frequent adverse events among participants in the ezogabine arms were fatigue and dizziness

¹Wainger BJ, Macklin EA, Vucic S, et al. *JAMA Neurol.* 2021;78(2):186–196. doi:10.1001/jamaneurol.2020.4300

²Error bars from JAMA paper represent standard deviations, results are statistically significant

PoM studies for ALS and epilepsy to support dose selection for PoC trials



SAD: Single Ascending Dose; MAD: Multiple Ascending Dose; PoM: Proof of Mechanism; PoC: Proof of Concept; PK/FE: Pharmacokinetics / Food Effect; HVs: Healthy Volunteers

ALS & epilepsy POM in healthy volunteers (QRL-101-05)

Evidence of both central & peripheral target engagement

• mNETT (Threshold Tracking) (ALS)

- Several measures were significantly altered by QRL-101 demonstrating inhibitory effects on peripheral nerve excitability
- SDTC, Rheobase, and several other measures appeared to have a dose dependent response

• Passive EEG (ALS / Epilepsy)

- Clear central activation/brain penetration with several EEG measures impacted
- Little to no activation of the slow wave frequency (delta and theta) associated with sedation and GABA-A activation

• TMS-EMG & TMS-EEG (ALS / Epilepsy)

- ICF TMS-EMG had statistically significant change with QRL-101 treatment
- Statistically significant changes to TMS Evoked Potential (TEP)

• Safety & tolerability

- Adverse event (AE) profile consistent with prior studies of QRL-101, no serious AEs or discontinuations due to AEs
- No clinically meaningful signal of somnolence or persistent dizziness has been observed in any study of QRL-101 to date
- Formulation work has resulted in lower cMax, while maintaining AUC, to alleviate GI AEs seen with liquid formulation

CONTEXT

Inhibition of nerve excitability

Target engagement & brain penetration

No sedative effects


Effect inhibition/ excitability ratio

Target engagement & brain penetration


QRL-101 is ~50% more potent than retigabine/Ezogabine on ALS biomarker SDTC in humans

QRL-101

Strength-Duration Time Constant (ms)



Strength-Duration Time Constant (%)



Ongoing studies will further support and confirm dosing for PoC studies

QRL-101

ALS PoM design (NCT06714396)

- Single-dose placebo-controlled design at three ascending dose levels
- 12 patients (four per dose level)
- Safety and tolerability in ALS patients
- PK/PD assessment at each dose level

Disease-relevant biomarkers collected

- Endpoints associated with peripheral nerve excitability threshold tracking
- Includes strength-duration, recovery cycle, threshold electrotonus and current / voltage; all output measures shown to be disrupted in ALS
- ALS biomarkers also included in previously reported healthy volunteer PoM (QRL-101-05)

Formulation & food effect (NCT06877624)



- 24 healthy volunteers
- Open label study to compare PK of three different formulations of QRL-101 (liquid formulation, two extended-release formulations)
- Select final formulation & dose for PoC studies

Study design & goals

- Initial treatment period to compare PK exposures in fasted healthy volunteers completed
- Three way cross-over to compare food effect and relative fed exposures of all three formulations
- Targeting lower cMax while maintaining AUC observed in liquid formulation studies, which demonstrated statistically significant effects on biomarkers predictive of efficacy in ALS and epilepsy

Topline data for both studies are expected Q2/Q3 2025

RNA STM



STMN2-targeting ASO leads RNA restoration franchise

STMN2 is the most consistently downregulated gene in sporadic ALS patients

- Restoration of STMN2 pre-mRNA mis-splicing through ASO treatment restores neuronal processes, Golgi outposts, and protects neuronal activity in human motor neurons with TDP-43 pathology
- Genetic target in sporadic ALS (90% of patients) and FTD (50% of patients) as well as Alzheimer's Disease (~33% of patients)
- Two approved ASO therapies for motor neuron diseases (Spinraza® for SMA and Qalsody® for ALS) demonstrate that an ASO therapy strategy to restore STMN2 in ALS patients is technologically feasible



- QurAlis is developing QRL-201, a highly potent splice-switching ASO targeting STMN2
- MAD study (ANQUR) expanded to dose range-finding portion (at two dose levels) and ongoing with favorable safety and tolerability profile to date
- Multiple biomarkers under assessment to support future development strategy
- QurAlis retains full global rights; CoM patent through 2039 plus potential PTE, pending issuance


Restoration of STMN2 levels by ASO is a promising therapeutic strategy for the majority of ALS patients

QRL-201 restores full-length STMN2 and protects neurons from degeneration phenotypes

Breakthrough discovery shows TDP-43 driven neurodegenerative phenotypes caused by STMN2 loss

Loss of TDP-43 from the nucleus (ALS hallmark) leads to loss of STMN2


TDP-43 loss causes loss of axons; rescue by restoring STMN2 levels




1. Melamed Z, López-Erauskin J, Baughn MW, et al. Premature polyadenylation-mediated loss of stathmin-2 is a hallmark of TDP-43-dependent neurodegeneration. *Nat Neurosci*. 2019;22(2):180-190. doi:10.1038/s41593-018-0293-z
2. Klim, J.R., Williams, L.A., Limone, F. et al. ALS-implicated protein TDP-43 sustains levels of STMN2, a mediator of motor neuron growth and repair. *Nat Neurosci* 22, 167–179 (2019). <https://doi.org/10.1038/s41593-018-0300-4>
3. Baughn, M. W., et al. (2023). "Mechanism of STMN2 cryptic splice-polyadenylation and its correction for TDP-43 proteinopathies." *Science* 379(6637): 1140-1149.
4. Krus et al., 2022 *Cell Rep* Jun 28;39(13):111001. doi: 10.1016/j.celrep.2022.111001

STMN2 is downregulated in ALS patients


Highley et al 2014 Microarray Laser Capture Motor neuron




Loss of STMN2 leads to denervation of muscles as measured by CMAP Amplitude



QRL-201 restores STMN2



QRL-201 restores neuronal processes




ANQUR study progressed to dose range-finding phase, focused on signal detection and safety

Enrollment initiated, interim data¹ expected H1 2026

QRL-201

AnQur




Design

Randomized, double-blind, placebo-controlled



Population

32 sporadic ALS patients
16 *C9orf72* ALS patients



Key entry criteria

Symptom onset within 24 months of screening
Slow vital capacity >50%
Clinical evidence of low motor neuron involvement
Stable dose of current treatment during study



Endpoints

1°: Safety & tolerability
2°: Efficacy & biomarker panel

¹ Exact interim data cut to be finalized

² *C9orf72* patients are a homogenous population with consistently decreased *STMN2* levels


Combination of clinical readouts and extensive biomarker analysis to inform optimal registrational studies

QRL-201

AnQur


Target engagement

- STMN2 levels




Biomarkers of neuronal loss

- NfL and other exploratory biomarkers



Mechanism of action (MOA)

- Motor excitability recordings (CMAP, M-Scan)
- Established NMJ innervation measurements (STMN2 MOA / efficacy)



Clinical measurements


- ALSFRS-R, ROADS, SVC, HHD
- Ventilation assistance-free survival
- Time-to-event measures




CMAP is a powerful disease progression and efficacy biomarker that measures muscle innervation in ALS

QRL-201


CMAP amplitude measures signal strength from nerve to muscle: number of innervating fibers




CMAP is used to monitor disease progression in ALS patients



CMAP amplitude predicts survival in ALS



CMAP amplitude can show therapeutic effect of ASO




1. Clarissa Crone, Christian Krarup, in *Handbook of Clinical Neurology*, 2013
2. Maathuis et al., 2013 *Amyotrophic Lateral Sclerosis and Frontotemporal Degeneration*, 2013; 14: 217–223
3. Yu et al., 2021 *Front Neurol*. 2021 Feb 11;12:574919. doi: 10.3389/fneur.2021.574919
4. McCampbell et al., 2018 *J Clin Invest*. 2018;128(8):3558–3567. <https://doi.org/10.1172/JCI99081>.
5. Krus et al., 2022 *Cell Rep* Jun 28;39(13):111001. doi: 10.1016/j.celrep.2022.111001


QRL-201 key take-aways



STMN2 is the most consistently observed mis-spliced protein in sporadic ALS leading to loss of function



Downregulation and loss of STMN2 alone in mice leads to loss of muscle innervation, motor neuron axonopathy and muscle atrophy, all hallmarks of ALS



QRL-201 restores STMN2 levels in human ALS motor neurons



Preliminary cohorts in the ANQUR study demonstrate that QRL-201 can be well tolerated in ALS patients at exposures far above the predicted minimally efficacious exposure



A dose range-finding study for QRL-201 is currently active in six countries with biomarkers that can measure efficacy in ALS patients

ANQUR study interim efficacy marker & safety data expected H1 2026
Next Ph2 / Ph3 study is a potential registrational study

RNA Restoration: FlexASO[®] Platform




Flex ASO[®] is the leading splice-modulation platform to restore RNA

FlexASO[®] is a proprietary ASO splice-modulator platform that incorporates a unique backbone, providing advantages over traditional ASOs

Attributes	Flex ASO	Traditional ASO
Size	✓	✓
Therapeutic Effect	✓ ✓	✓
Toxicity	✓ ✓	✓
CMC	✓	✓
Distribution	✓ ✓	Known for spinal cord and frontal cortex

Potential to overcome modality-specific, dose-limiting toxicities

FlexASO[®] demonstrates statistically significant increase in RNA restoration vs. parent



Genetic understanding of diseases and tech breakthroughs generate opportunity to develop RNA restoration therapies

Technological breakthroughs



Patient-derived human disease models increase probability of success



Technology to potentially support 2x / year dosing has been developed (e.g., FlexASO® tech.)



Technology to cross blood-brain-barrier has matured (e.g., Transferrin receptor)

Disease-modifying RNA therapies

RNA restoration oligonucleotides targets

Diseases caused by mis-splicing (e.g., SMA, ALS, FTD, Fragile X, PSP)



Diseases caused by haploinsufficiency (e.g., Dravet)



Previously undruggable targets for large indications



Driving scientific breakthroughs into powerful precision medicines




Groundbreaking science

Next-gen precision medicines developed by **human disease models** pioneered by QurAlis founders

Targeting **RNA restoration** in validated genetic disease resulting from **mis-splicing targets** in neurodegeneration and beyond

Utilizing **biomarkers** for patient selection, target engagement, and efficacy



First & best-in-class programs

Multiple programs in the clinic with near-term data readouts

Three first-in-class programs for sporadic ALS **on novel genetic targets**

Proprietary FlexASO® platform can potential expand to additional RNA restoration therapies



World-class team to execute

Experienced executive team

UNC13A partnership with Lilly underscores value of FlexASO® platform

\$143.5M equity raised, in addition to Lilly partnership upfront



QuralisTM

Thank you

For more information contact:
Kasper Roet (CEO & Cofounder)
kasper.roet@quralis.com

Jason Brown (CFO)
jason.brown@quralis.com